

570

NET WT. 50 LBS. (22.67 kg)

570

**Burkman  
Feeds**

**25% CAPRINE BALANCER-  
RUM**

MEDICATED

**A MIXING SUPPLEMENT TO MAKE COMPLETE  
GOAT FEEDS**

For the prevention of coccidiosis caused by *Eimeria crandallis*,  
*Eimeria christenseni*, and *Eimeria ninakohlyakimovae*.

**ACTIVE DRUG INGREDIENT**

Monensin..... 60 g/ton

**GUARANTEED ANALYSIS**

Crude Protein .....	Minimum .....	25.00%
(This includes not more than 3.00% equivalent crude protein from non-protein nitrogen.)		
Crude Fat .....	Minimum .....	2.00%
Crude Fiber .....	Maximum .....	8.00%
Calcium .....	Minimum .....	1.90%
Calcium .....	Maximum .....	2.40%
Phosphorus .....	Minimum .....	0.85%
Salt .....	Minimum .....	1.25%
Salt .....	Maximum .....	1.75%
Copper.....	Minimum .....	12 ppm
Copper.....	Maximum .....	16 ppm
Selenium .....	Minimum .....	0.9 ppm
Vitamin A .....	Minimum IU/Lb.....	60,000
Vitamin D <sub>3</sub> .....	Minimum IU/Lb.....	10,000
Vitamin E .....	Minimum IU/Lb.....	60

**INGREDIENTS**

Plant protein products, processed grain by-products, forage products, molasses products, ammonium chloride, calcium carbonate, magnesium sulfate, potassium sulfate, salt, zinc sulfate, zinc methionine complex, manganese sulfate, ethylenediamine dihydriodide, cobalt carbonate, dried yeast fermentation solubles, selenium yeast, brewer's dried yeast, vitamin A supplement, vitamin D<sub>3</sub> supplement, vitamin E supplement, yeast culture.

**IMPORTANT: SEE REVERSE SIDE OF TAG FOR FEEDING DIRECTIONS AND CAUTION STATEMENTS.**

Manufactured By

**BURKMANN FEEDS**

**Danville, Kentucky 40422  
NET WT. 50 LBS. (22.67 kg)**

**Burkman  
Feeds**

**25% CAPRINE BALANCER-  
RUM**

MEDICATED

**A MIXING SUPPLEMENT TO MAKE COMPLETE  
GOAT FEEDS**

For the prevention of coccidiosis caused by *Eimeria crandallis*,  
*Eimeria christenseni*, and *Eimeria ninakohlyakimovae*.

**ACTIVE DRUG INGREDIENT**

Monensin..... 60 g/ton

**GUARANTEED ANALYSIS**

Crude Protein .....	Minimum .....	25.00%
(This includes not more than 3.00% equivalent crude protein from non-protein nitrogen.)		
Crude Fat .....	Minimum .....	2.00%
Crude Fiber.....	Maximum .....	8.00%
Calcium .....	Minimum .....	1.90%
Calcium .....	Maximum .....	2.40%
Phosphorus .....	Minimum .....	0.85%
Salt .....	Minimum .....	1.25%
Salt .....	Maximum .....	1.75%
Copper.....	Minimum .....	12 ppm
Copper.....	Maximum .....	16 ppm
Selenium .....	Minimum .....	0.9 ppm
Vitamin A .....	Minimum IU/Lb.....	60,000
Vitamin D <sub>3</sub> .....	Minimum IU/Lb.....	10,000
Vitamin E .....	Minimum IU/Lb.....	60

**INGREDIENTS**

Plant protein products, processed grain by-products, forage products, molasses products, ammonium chloride, calcium carbonate, magnesium sulfate, potassium sulfate, salt, zinc sulfate, zinc methionine complex, manganese sulfate, ethylenediamine dihydriodide, cobalt carbonate, dried yeast fermentation solubles, selenium yeast, brewer's dried yeast, vitamin A supplement, vitamin D<sub>3</sub> supplement, vitamin E supplement, yeast culture.

**IMPORTANT: SEE REVERSE SIDE OF TAG FOR FEEDING DIRECTIONS AND CAUTION STATEMENTS.**

Manufactured By

**BURKMANN FEEDS**

**Danville, Kentucky 40422**

**Burkman  
Feeds**

**25% CAPRINE BALANCER-  
RUM**

MEDICATED

**A MIXING SUPPLEMENT TO MAKE COMPLETE  
GOAT FEEDS**

For the prevention of coccidiosis caused by *Eimeria crandallis*,  
*Eimeria christenseni*, and *Eimeria ninakohlyakimovae*.

**ACTIVE DRUG INGREDIENT**

Monensin..... 60 g/ton

**GUARANTEED ANALYSIS**

Crude Protein .....	Minimum .....	25.00%
(This includes not more than 3.00% equivalent crude protein from non-protein nitrogen.)		
Crude Fat .....	Minimum .....	2.00%
Crude Fiber.....	Maximum .....	8.00%
Calcium.....	Minimum .....	1.90%
Calcium.....	Maximum .....	2.40%
Phosphorus .....	Minimum .....	0.85%
Salt .....	Minimum .....	1.25%
Salt .....	Maximum .....	1.75%
Copper.....	Minimum .....	12 ppm
Copper.....	Maximum .....	16 ppm
Selenium.....	Minimum .....	0.9 ppm
Vitamin A .....	Minimum IU/Lb.....	60,000
Vitamin D <sub>3</sub> .....	Minimum IU/Lb.....	10,000
Vitamin E .....	Minimum IU/Lb.....	60

**INGREDIENTS**

Plant protein products, processed grain by-products, forage products, molasses products, ammonium chloride, calcium carbonate, magnesium sulfate, potassium sulfate, salt, zinc sulfate, zinc methionine complex, manganese sulfate, ethylenediamine dihydriodide, cobalt carbonate, dried yeast fermentation solubles, selenium yeast, brewer's dried yeast, vitamin A supplement, vitamin D<sub>3</sub> supplement, vitamin E supplement, yeast culture.

**IMPORTANT: SEE REVERSE SIDE OF TAG FOR FEEDING DIRECTIONS AND CAUTION STATEMENTS.**

Manufactured By

**BURKMANN FEEDS**  
Danville, Kentucky 40422  
NET WT. 50 LBS. (22.67 kg)

**570**

**PRODUCT NUMBER 570**

**FEEDING DIRECTIONS**

**BURKMANN 25% CAPRINE BALANCER-RUM** is intended for mixing with soybean hulls to make a complete feed for growing goats. Mix 660 pounds of **BURKMANN 25% CAPRINE BALANCER-RUM** with 1340 pounds of soybean hulls to make a ton of complete feed. On a small batch basis this is the equivalent of mixing ½ pound of **BURKMANN 25% CAPRINE BALANCER-RUM** with 1.0 pound of soybean hulls. The final feed mixture will contain 20 grams/ton of Rumensin and should be fed to growing goats as the sole grain ration. For hand feeding, insure that adequate trough space is provided for all goats to eat at once. Provide **BURKMANN GOAT MINERAL** and clean, fresh water free choice at all times.

**CAUTION: Do not allow horses or other equines access to feeds containing Monensin. Ingestion of Monensin by horses has been fatal. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal in cattle and could be fatal in goats. Monensin medicated feed is intended for use in cattle or in goats. Consumption by unapproved species may result in toxic reaction. Do not feed to lactating goats.**

04076

**PRODUCT NUMBER 570**

**FEEDING DIRECTIONS**

**BURKMANN 25% CAPRINE BALANCER-RUM** is intended for mixing with soybean hulls to make a complete feed for growing goats. Mix 660 pounds of **BURKMANN 25% CAPRINE BALANCER-RUM** with 1340 pounds of soybean hulls to make a ton of complete feed. On a small batch basis this is the equivalent of mixing ½ pound of **BURKMANN 25% CAPRINE BALANCER-RUM** with 1.0 pound of soybean hulls. The final feed mixture will contain 20 grams/ton of Rumensin and should be fed to growing goats as the sole grain ration. For hand feeding, insure that adequate trough space is provided for all goats to eat at once. Provide **BURKMANN GOAT MINERAL** and clean, fresh water free choice at all times.

**CAUTION: Do not allow horses or other equines access to feeds containing Monensin. Ingestion of Monensin by horses has been fatal. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal in cattle and could be fatal in goats. Monensin medicated feed is intended for use in cattle or in goats. Consumption by unapproved species may result in toxic reaction. Do not feed to lactating goats.**

04076

**PRODUCT NUMBER 570**

**FEEDING DIRECTIONS**

**BURKMANN 25% CAPRINE BALANCER-RUM** is intended for mixing with soybean hulls to make a complete feed for growing goats. Mix 660 pounds of **BURKMANN 25% CAPRINE BALANCER-RUM** with 1340 pounds of soybean hulls to make a ton of complete feed. On a small batch basis this is the equivalent of mixing ½ pound of **BURKMANN 25% CAPRINE BALANCER-RUM** with 1.0 pound of soybean hulls. The final feed mixture will contain 20 grams/ton of Rumensin and should be fed to growing goats as the sole grain ration. For hand feeding, insure that adequate trough space is provided for all goats to eat at once. Provide **BURKMANN GOAT MINERAL** and clean, fresh water free choice at all times.

**CAUTION: Do not allow horses or other equines access to feeds containing Monensin. Ingestion of Monensin by horses has been fatal. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal in cattle and could be fatal in goats. Monensin medicated feed is intended for use in cattle or in goats. Consumption by unapproved species may result in toxic reaction. Do not feed to lactating goats.**

04076